



## Small-scale polygenetic mélanges in the Ligurian accretionary complex, Northern Apennines, Italy, and the role of shale diapirism in superposed mélange evolution in orogenic belts

Giulia Codegone <sup>a,\*</sup>, Andrea Festa <sup>a</sup>, Yildirim Dilek <sup>b</sup>, Gian Andrea Pini <sup>c</sup>

<sup>a</sup> Dipartimento di Scienze della Terra, Università di Torino, 10125 Torino, Italy

<sup>b</sup> Department of Geology and Environmental Earth Science, Miami University, Oxford, OH 45056, USA

<sup>c</sup> Dipartimento di Scienze della Terra e Geologico-Ambientali, Università di Bologna, 40127 Bologna, Italy

### ARTICLE INFO

#### Article history:

Received 27 March 2011

Received in revised form 28 January 2012

Accepted 3 February 2012

Available online 11 February 2012

#### Keywords:

Polygenetic mélanges

Broken formation

Tectonic and sedimentary mélanges

Accretionary wedge

Shale diapirism

Northern Apennines

### ABSTRACT

The *Argille varicolori* (Varicolored scaly clays) of the External Ligurian Units of the Northern Apennines have been widely described as a typical unmetamorphosed broken formation (i.e., a chaotic unit without exotic blocks), produced by offscraping and tectonic imbrication during the evolution of the Ligurian accretionary wedge. Geological mapping and integrated structural and stratigraphic observations show that the *Argille varicolori* consist of diverse types of small-scale mélanges (non-mappable at a 1:25,000 scale) forming a composite chaotic unit, in which the superposition of tectonic, sedimentary and diapiric processes resulted in the occurrence of polygenetic chaotic bodies at different scales. These mélange units record the evolution of the Ligurian accretionary wedge from subduction to collision and intracontinental deformation. Tectonically Disrupted Body 1 (TDB1) comprises boudinage and pinch-and-swell structures formed by layer-parallel extension/contraction at the wedge front of the Ligurian accretionary complex during the late Cretaceous–middle Eocene. It is interleaved with non-mappable Gravity-driven Chaotic Bodies (GCB) developed during alternating episodes of accretion and removal of material at the wedge front. The late Oligocene–early Miocene out-of-sequence thrusting related to the collisional episodes in the Apennines overprinted the previously formed chaotic bodies and formed a polygenetic tectonic mélange (Tectonically Disrupted Body 2, TDB2). This unit is characterized by a structurally ordered block-in-matrix fabric and by the gradual decrease of stratal disruption away from the regional thrust. Overpressurized fluids concentrated along the shear surfaces, and the scaly cleavage planes facilitated the diapiric upward movement of unconsolidated sediments in the early Miocene. This process produced non-mappable shale dike injections (DDB1) and mappable Diapirically Disrupted Bodies (DDB2), which show an internal zonation of deformation. This deformation reworked the previously formed chaotic bodies. Although some of these polygenetic mélanges cannot be mapped at a 1:25,000 scale, their careful documentation provides a better understanding of time-progressive, scale-independent mélange-forming processes.

© 2012 Elsevier B.V. All rights reserved.

### 1. Introduction

Accretionary complexes in convergent margins constitute a primary tectonic setting for the formation of chaotic bodies of mixed rocks and mélanges. Comparative studies of modern and ancient accretionary complexes are, therefore, a natural laboratory to investigate the mode and nature of all geological processes involved in mélange formation and their mechanisms and interplay (Ogawa et al., 2011).

Tectonic, sedimentary and diapiric processes that play the most important role in the formation of chaotic bodies and mélanges

occur at all scales in convergent margin settings. While it is possible to decipher the internal architecture and large-scale structures in modern accretionary complexes by indirect methods of observations (i.e., seismic imaging), it is difficult to do so when the scale of observation is smaller (mesoscale to outcrop scale). Submersible investigations and drill cores provide *in situ* samples and measurements from accretionary complexes (Anma et al., 2010, 2011; Kawamura et al., 2009; Ogawa et al., 2011), but these methods are usually highly expensive and limited in coverage. The well-preserved on-land examples of ancient accretionary complexes are, therefore, highly important to conduct three-dimensional studies of chaotic bodies of mixed rocks and mélanges at various scales, and to better document different processes and their superposition during progressive evolution of accretionary complexes (Aalto, 1981; Cowan, 1985; Cowan and Pini, 2001; Festa et al., 2010a, 2010b; Ghikas et al., 2010;

\* Corresponding author. Tel.: +39 347 0448914.

E-mail address: [giulia.code@gmail.com](mailto:giulia.code@gmail.com) (G. Codegone).